

AMENDMENTS TO THE CLAIMS

1. (Previously Presented) A method for inducing a mucosal immune response, comprising:
administering to a subject in need of a mucosal immune response an effective amount for
inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a
sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, and

an antigen,

wherein the antigen is not encoded in a nucleic acid vector, the oligonucleotide and the
antigen are both administered vaginally, rectally, intranasally, ocularly, or by inhalation to the
subject, a cytokine and an immune stimulating complex are not administered to the subject, and the
antigen is not a *Streptococcus pneumoniae* antigen.

2.-3. (Cancelled)

4. (Previously Presented) The method of claim 1, wherein the antigen is administered
concurrently with the oligonucleotide.

5. (Previously Presented) The method of claim 1, wherein the antigen is delivered in
conjunction with a colloidal dispersion system.

6. (Original) The method of claim 5, wherein the colloidal dispersion system is selected
from the group consisting of macromolecular complexes, nanocapsules, microspheres, beads, and
lipid-based systems.

7. (Original) The method of claim 6, wherein the lipid-based system is selected from the
group consisting of oil-in-water emulsions, micelles, mixed micelles, and liposomes.

8. (Previously Presented) The method of claim 1, further comprising the step of administering a non-oligonucleotide mucosal adjuvant in conjunction with the antigen.

9. (Previously Presented) The method of claim 8, wherein the non-oligonucleotide mucosal adjuvant is selected from the group consisting of cholera toxin, derivatives of cholera toxin, heat-labile enterotoxin, derivatives of heat-labile enterotoxin, alum, monophosphoryl lipid A (MLP), muramyl dipeptide (MDP), saponins, QS21, cytokines, oil-in-water and other emulsion formulations, squalene-in-water emulsion stabilized with Span 85 and Tween 80 (MF59), syntex adjuvant formulation (SAF), Montanide ISA 720 and oil-in-water emulsion containing stabilizing detergent and micelle-forming agent and poly (PCPP) polymers.

10.-11. (Cancelled)

12. (Previously Presented) The method of claim 1, wherein the subject is a subject at risk of developing an infectious disease.

13. (Previously Presented) The method of claim 1, wherein the subject is at risk of developing cancer.

14. (Cancelled)


15. (Original) The method of claim 1, wherein the oligonucleotide includes a phosphate backbone modification which is a phosphorothioate or phosphorodithioate modification.

16. (Original) The method of claim 15, wherein the phosphate backbone modification occurs at the 5' end of the oligonucleotide.

17. (Original) The method of claim 15, wherein the phosphate backbone modification occurs at the 3' end of the oligonucleotide.

18. (Original) The method of claim 1, wherein X_1X_2 are nucleotides selected from the group consisting of: GpT, GpG, GpA, ApA, ApT, ApG, CpT, CpA, CpG, TpA, TpT, and TpG; and X_3X_4 are nucleotides selected from the group consisting of: TpT, CpT, ApT, TpG, ApG, CpG, TpC, ApC, CpC, TpA, ApA, and CpA.

19. (Original) The method of claim 1, wherein the oligonucleotide has a sequence including at least the following formula:



wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, N is a nucleic acid sequence composed of from about 0-25 nucleotides.

20. (Previously Presented) The method of claim 1, wherein the antigen is selected from the group consisting of cells, cell extracts, proteins, polypeptides, peptides, polysaccharides, polysaccharide conjugates, peptide mimics of polysaccharides, lipids, glycolipids, carbohydrates, viruses and viral extracts and parasites.

21. (Cancelled)

22. (Previously Presented) The method of claim 1, wherein the antigen is obtained from an infectious organism selected from the group consisting of infectious bacteria, infectious viruses, infectious parasites, and infectious fungi.

23-24. (Cancelled)

25. (Previously Presented) The method of claim 1, further comprising administering a B-7 costimulatory molecule.

26. (Previously Presented) The method of claim 1, wherein the mucosal immune response is induced in a remote site.

27. (Original) The method of claim 1, further comprising administering a boost of the oligonucleotide.

28. (Original) The method of claim 8, further comprising administering a boost of the oligonucleotide and the non-oligonucleotide mucosal adjuvant.

29-128. (Cancelled)

129. (Previously Presented) The method of claim 1, further comprising identifying a subject in need of a mucosal immune response.

130.-134. (Cancelled)

135. (Previously Presented) The method of claim 1, wherein the antigen is a viral antigen.

136. (Previously Presented) A method for inducing a mucosal immune response, comprising:

administering to a subject in need of a mucosal immune response an effective amount for inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides,

a non-oligonucleotide mucosal adjuvant that is not an immune stimulating complex, and an antigen,

wherein the antigen is not encoded in a nucleic acid vector, and wherein the oligonucleotide, the antigen, and the non-oligonucleotide mucosal adjuvant are all administered rectally, intravaginally, or ocularly, to the subject, and a cytokine is not administered to the subject.

137. (Previously Presented) A method for inducing a mucosal immune response, comprising:

administering to a subject in need of a mucosal immune response an effective amount for inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, and
a viral antigen,

wherein the antigen is not encoded in a nucleic acid vector, the oligonucleotide and the antigen are both administered vaginally, rectally, intranasally, ocularly, or by inhalation to the subject, and a cytokine and an immune stimulating complex are not administered to the subject.

138. (Previously Presented) A method for inducing a mucosal immune response, comprising:

administering to a subject in need of a mucosal immune response an effective amount for inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, and

passively exposing the subject to an antigen,

wherein the antigen is not encoded in a nucleic acid vector, oligonucleotide administration and antigen exposure both occur vaginally, rectally, intranasally, or by inhalation, and a cytokine and an immune stimulating complex are not administered to the subject.

139. (Previously Presented) A method for inducing a mucosal immune response, comprising:

administering to a subject in need of a mucosal immune response an effective amount for inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, and
an antigen,

wherein the antigen is not encoded in a nucleic acid vector, the oligonucleotide and the antigen are both administered vaginally, rectally, or ocularly to the subject, and a cytokine and an immune stimulating complex are not administered to the subject.

140. (Previously Presented) The method of claim 139, wherein the antigen is a viral antigen.

141. (Previously Presented) A method for inducing a mucosal immune response, comprising:

administering to a subject in need of a mucosal immune response an effective amount for inducing a mucosal immune response of an oligonucleotide 8 to 100 nucleotides in length, having a sequence including at least the following formula:



wherein C is unmethylated, wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, and
an antigen,

wherein the antigen is not encoded in a nucleic acid vector and is not a *Streptococcus pneumoniae* antigen, the oligonucleotide and the antigen are both administered intranasally or by inhalation to the subject, and a cytokine and an immune stimulating complex are not administered to the subject.

142. (Previously Presented) The method of claim 136, wherein the antigen is selected from the group consisting of cells, cell extracts, proteins, polypeptides, peptides, polysaccharides,

polysaccharide conjugates, peptide mimics of polysaccharides, lipids, glycolipids, carbohydrates, viruses and viral extracts and parasites.

143. (Cancelled)

144. (Previously Presented) The method of claim 138, wherein the antigen is selected from the group consisting of cells, cell extracts, proteins, polypeptides, peptides, polysaccharides, polysaccharide conjugates, peptide mimics of polysaccharides, lipids, glycolipids, carbohydrates, viruses and viral extracts and parasites.

145. (Previously Presented) The method of claim 139, wherein the antigen is selected from the group consisting of cells, cell extracts, proteins, polypeptides, peptides, polysaccharides, polysaccharide conjugates, peptide mimics of polysaccharides, lipids, glycolipids, carbohydrates, viruses and viral extracts and parasites.

146. (Previously Presented) The method of claim 141, wherein the antigen is selected from the group consisting of cells, cell extracts, proteins, polypeptides, peptides, polysaccharides, polysaccharide conjugates, peptide mimics of polysaccharides, lipids, glycolipids, carbohydrates, viruses and viral extracts and parasites.